Lumbar Puncture / Myelogram Questionnaire

Inpatient	Outpatient			
Must have head study CT or MRI (unless waived by RADIOLOGIST) Head study reviewed by radiologist/ RA? Yes No	Are you on blood thinners?			
Is the patient on blood thinners? Yes No Medication Last Dose	If yes, instruct patient to call provider and confirm anticoagulation medication can be withheld for the recommended amount of time.			
If yes, follow anticoagulation rules and PLT/INR to be done prior to exam	Are you able to lay on your stomach?			
PLT INR >50,000 < 1.5	Do you have a ride for your exam? (MUST HAVE DRIVER) Yes No			
Is patient able to lay prone?	Are you allergic to chlorohexidine or lidocaine? Yes No			
Is patient allergic to chlorohexidine or lidocaine? Yes No	Myelogram pts: Do you have an lodine or contrast allergy? Yes No			
Is patient well hydrated? Yes No If no, PLEASE HYDRATE	If yes, pt. must have 13 hour prep			
Myelogram pts:	If patient is unable to consent, POA must provide consent at the time of the exam			
lodine or contrast allergy? Yes No	Chemo injections: 1 day prior to exam- call pharmacy/fax order (verify received),			
If yes, pt must have 13 hour prep	inform RCU of exam, RCU nurse to obtain chemo and handoff to RAD			
Nurse to begin prep at for exam @	Appointment time: Check in time:			
Is the patient able to consent? Yes No				
If no, will a family member/ POA be available to provide consent?				
Transport: Pt needs to travel in bed/stretcher, send stickers with patient	**Remind patient to hydrate before exam & limit activity post exam**			
Intrathecal chemo injections: pt's nurse must bring chemo meds and stay for exam (double check chemo meds ordered and available for scheduled exam)				

	Pradaxa (Dabigatran) Off 4	Therapeutic Lovenox	Prophylactic	Arixtra	Plavix (Clopidogrel)	ASA (Aspirin)	NSAIDs
MEDICATION	days in ESRD patients	(Enoxaparin) 1mg/kg BID	Lovenox	(Fondaparinux)	Pletal (Cilostazol)	Aggrenox	
			(30-40 mg daily)	Xarelto	Effient (Prasugrel)	(Aspirin	
	Coumadin (Warfarin)	or		(Rivaroxiban)	Brilinta (Ticagrelor)	Dipyridamole)	
	CHECK INR IF		SQ heparin	Eliquis (Apixaban)	Ticlid (Ticlopidine)		
	COUMADIN HELD	IV heparin: Consider	(5000 U BID-TID)	Fragmin (Dalteparin)			
		holding 2-4 hours if aPTT					
PROCEDURE	Do not give FFP for elevated	> 1.5x upper limits of					
	INR in patients with stable	hospital lab's nomogram.					
	chronic liver disease						
Lumbar Drains	Off 3 days.	Hold 24 hours	Hold 12 hours pre	Off 24 hours	Off 5 days,	Continue	Continue
Lumbar Puncture	INR should be 1.5 or less	pre and post	and 24 hours post	pre and post	if possible		
Myelograms							